arXiv:0910.2355v2 [hep-ph] 7 Dec 2009

Quark-hadron duality in lepton scattering off nucleons

Krzysztof M. Graczyk

Institute of Theoretical Physics, Wrocław University, pl. M. Borna 9, 50-204, Wrocław, Poland

Abstract. Quark-hadron (QH) duality in lepton scattering off nucleons is studied with the resonance quark model. It is shown that in the case of neutrino scattering off an isoscalar target the duality is simultaneously observed for charged and neutral currents $xF_1^{\nu N}$, $F_2^{\nu N}$, and $xF_3^{\nu N}$ weak structure functions.

We demonstrate that the QH duality can be useful property for modeling structure functions in the so-called resonance region. As an example it is shown that combining relativistic quark model predictions with duality arguments allows a construction of the inclusive resonance F_2^{ep} structure function.

Keywords: Bloom-Gilman duality, nucleon structure functions, weak single pion production
PACS: 25.30.Pt, 13.15.+g

INTRODUCTION

Historically *quark-hadron duality* was discovered as a remarkable relationship between hadronic scattering amplitudes in s (low energy scale interaction) and t (high energy scale interaction) channels.

The realization of the QH duality in inelastic electron-proton scattering was discovered by Bloom and Gilman [1] (for review see [2]). It was observed that the ep inelastic νW_2 resonance data scales as the deep inelastic scattering (DIS) one. In practise, it was shown that the resonance $\nu W_2^{ep} = F_2^{ep}$ data, considered as a function of $\omega' = 1 + W^2/Q^2$ scaling variable, is averaged by the DIS scaling curve.

The *Bloom-Gilman (BG) duality* constitutes a kind of equivalence between the resonance and DIS structure functions. As was obtained by De Rujula *et al.* [3] the duality phenomenon can be explained by doing the twist expansion of the F_2^{ep} structure function. In low-Q^2, the leading twist of the F_2^{ep} dominates, while in high-Q^2 higher twist correction become more relevant. Thus the appearance of the BG duality means the suppression of the contribution from higher twists. De Rujula *et al.* showed also that the Nachtman variable $\xi(x,Q^2) = 2x/\left(1+\sqrt{1+4x^2M^2/Q^2}\right)$ is a correct scaling variable to discuss QCD scaling violations, and consequently the proper scaling variable to study the BG duality.

From the practical point of view the duality means equality between two integrals:

$$\int_{\xi_{min}}^{\xi_{max}} d\xi F^{res}(\xi, Q^2_{res}) \approx \int_{\xi_{min}}^{\xi_{max}} d\xi F^{DIS}(\xi, Q^2_{DIS}), \quad (1)$$

where F^{res} and F^{DIS} denote the structure functions at resonance and DIS regions respectively. The ξ_{min}, ξ_{max} are defined by the limits of resonance region, namely:

$$\xi_{min/max} = \xi(Q^2_{res}, W_{max/min}). \quad (2)$$

For the lower bound of the resonance region we take $W_{min} = M + m_\pi$, while W_{max} will vary from 1.6 to 2.2 GeV. The typical four-momentum transfer in the resonance region for ep scattering is $0 < Q^2_{res} < 5$ GeV^2. For inclusive νN scattering we consider $0 < Q^2_{res} 3$ GeV^2. The characteristic four-momentum transfer for DIS is $Q^2_{DIS} \sim 10$ GeV^2. With such kinematical settings F^{DIS} is integrated over a different kinematical domain, in invariant, mass than F^{res}.

Recently the BG duality has been carefully studied by Niculescu *et al.* [4]. The new precision resonance JLab data has been discussed together with the old measurements. The BG duality was quantitatively studied in a wide kinematical region: $0.3 < Q^2_{res} < 5$ GeV^2, $1.1 < W^2 < 4$ GeV^2. The results of this analysis confirmed the appearance of the BG duality but with accuracy depending on the prominent resonance regions.

The duality is also observed in electron-deuteron scattering [5], which is also evidence for the duality between electromagnetic neutron structure functions.

It seems that analogously as for the eN scattering, the duality should be observed also for weak nucleon structure functions, which are used to describe the inclusive νN interaction. In contrast to the eN scattering the experimental evidence for the duality in νN scattering does not exist. The neutrino scattering data, collected in the resonance region, is still too imprecise to order to extract directly the resonance structure functions. On the other hand the knowledge of the DIS structure functions for inclusive νN scattering is also limited (see [6]).

Given the lack of appropriate neutrino scattering data, the duality in νN scattering has been studied only with the phenomenological descriptions [7, 8, 9], which have



FIGURE 1. In the left panel, plots of F_2^{ep} structure functions are presented. The quark model predictions (solid line) are compared with the inclusive CLAS data [18] (full squares and triangles and open circles) and DIS scaling curve (dashed line). In the right panel the plots of the ratios (4), computed for W_{max} =1.6, 1.8, 2.0 and 2.0 GeV, are shown.

been fine tuned to the old neutrino-deuteron bubble chamber scattering data.

Assuming that the duality is a property of the electromagnetic and weak nucleon structure functions can lead to the additional constraint on: the parameterization of the nucleon and resonance form factors[1] as well as the nucleon inclusive structure functions.

This talk presents an extension of the analysis published in Ref. [8]. The theoretical framework is formulated in Section 2. In Section 3 we demonstrate that combining quark model predictions and duality arguments leads to the reasonable parametrization of the inclusive F_2^{ep} structure function in the resonance region. The second aim of the talk is to investigate the BG duality in neutrino scattering off nucleons. In the last section we shortly discuss the duality in lepton scattering off nuclei.

THEORETICAL FRAMEWORK

In order to compute the resonance structure functions we apply an updated [11] Rein-Sehgal (RS) model [12]. The RS model [12] was devoted to describe the 1π production induced by neutrino-nucleon interaction in a wide kinematical region (up to $W = 2.0$ GeV). The description is based on the relativistic harmonic oscillator quark model [13]. Even thought the RS model is relatively simple and old it is still the reference description for the experimental data analysis in modern experiments like K2K and MiniBooNE.

In the RS model the vector and axial hadronic currents are described with the nucleon vector and axial elastic form factors but, as it was shown in Ref. [11], applying these form factors leads to the underestimation of the vector part and overestimation of the axial contribution. In this analysis we use the vector and axial form factors proposed in Ref. [11]. The updated vector contribution agrees with the recent fits of the resonance-like electroproduction data (in the $P_{33}(1232)$ resonance region), while the new axial form factor was obtained as a simultaneous fit to the ANL and BNL 1π production data[2]. The updated RS model works efficiently and fits very well to the recent MiniBooNE measurements [14].

The mechanism for the duality appearance in lepton scattering off nucleons is partially explained by quark model arguments [15]. The RS model as an example of a quark model with strong experimental applications seems to be an interesting framework for phenomenological studies of the duality in νN scattering.

Close & Isgur claimed that one of the necessary conditions for the realization of the duality is taking into consideration at least one complete set of resonances of each symmetry-type. Thus, for completeness of our analysis we consider the following list of resonances[3]: $P_{33}(1232), P_{11}(1440), D_{13}(1520), S_{11}(1535), S_{31}(1620),$ $S_{11}(1650),$ $P_{33}(1600),$ $D_{13}(1700),$ $D_{15}(1675),$ $F_{15}(1680), D_{33}(1700), P_{11}(1710), P_{13}(1720), P_{31}(1750),$

---

[1] In Ref. [3] local duality in elastic *ep* scattering is discussed. The applications can be found in Ref. [10].

[2] The axial form factor is given by Eq. 52 from the erratum of Ref. [11].

[3] The interference terms between resonances are also included in the calculus.



FIGURE 2. In the left panel the predictions of the resonance structure function computed with Eq. 6 (thin solid line) are shown. We plot also the scaling function (thick dashed line), valence scaling function (thin dashed line) and resonance fit, given by Eq. 9 of Ref. [4], (dotted line). In the right panel the plots of the ratios Eqs. 4) and 5, computed for 1.15<W<3.9 GeV^2, are shown (solid and dashed lines respectively). The results are compared with the data from [4].

$P_{13}(1900)$, $S_{31}(1900)$, $P_{31}(1910)$, $F_{35}(1905)$, $F_{37}(1950)$, $P_{33}(1920)$, $F_{17}(1990)$, $F_{15}(2000)$, $F_{35}(2000)$, $G_{17}(2190)$, $G_{19}(2250)$. However, one needs to remember that for larger values of the hadronic invariant mass the harmonic oscillator quark model predictions deviate from the experimental measurements. Therefore extending the number of resonances in the discussion can give only a hint on understanding the duality.

In order to quantitatively study the BG duality we introduce the ratio:

$$R_i\left(f_i, Q^2_{res}, Q^2_{DIS}\right) = \frac{\displaystyle\int_{\xi_{min}}^{\xi_{max}} d\xi\, f^{res}(\xi, Q^2_{res})}{\displaystyle\int_{\xi_{min}}^{\xi_{max}} d\xi\, f^{DIS}(\xi, Q^2_{DIS})}, \qquad (3)$$

where the limits of integrals are defined by (2), while f_i^{DIS} and f_i^{res} are computed with GRV94 PDF's [16] and the updated RS model respectively.

F_2^{ep} STRUCTURE FUNCTION

In order to test our theoretical model we compare the theoretical predictions for inclusive F_2^{ep} structure function with the resonance CLAS data [18]. As can be seen in Fig. 1 the quark model predictions are systematically below the data but it is not a surprise, because the non-resonant background (NoBG) contribution is not taken into account. On the other hand, the experimental data, and theoretical predictions seem to lie along the valence $F_2^{ep,val}$ curve (see Fig. 2). This property of the resonance data has been already studied in Ref. [5].

The quantitative verification of our approach is done by computing the ratio (see Fig. 1)

$$R_2^{ep}(Q^2_{res}) \equiv R\left(F_2^{ep}, Q^2_{res}, Q^2_{DIS} = 10\ \mathrm{GeV}^2\right), \qquad (4)$$

where we consider five different values of ξ_{min}, namely W_{max} =1.6, 1.8, 2 and 2.2 GeV. It is seen that, above $Q^2_{res} = 0.5$, the ratios (4) computed for $W_{max} = 2$ GeV and $W_{max} = 2.2$ GeV slowly vary on Q^2_{res} ($R_2^{ep}(Q^2_{res} > 0.5) \sim 0.6$ and $R_2^{ep}(Q^2_{res} > 0.5) \sim 0.7$ respectively). The weak Q^2_{res} dependence of $R_2^{ep}(Q^2_{res})$ is evidence for the same scaling behavior of the resonance predictions and the DIS structure function, however, it is clear that some strength is missing to get a perfect duality.

The scaling is broken below $Q^2_{res} = 0.5$ GeV^2, where $R_2^{ep}(Q^2_{res})$ rapidly tends to zero[4]. The argument which explains the mechanism of the duality violation (below $Q^2_{res} = 0.5$ GeV^2) was given in [17]: in the quark models the duality is strongly related to the dominance of the magnetic contribution, while at low Q^2, the electric and magnetic multipoles have comparable strengths.

In order to compare more qualitatively with the experimental measurements we compute the ratio (Eq. 4) for $1.1 < W^2 < 3.9$ GeV^2 and compare it with the analogous one obtained by Niculescu *et al.* [4] (here the DIS contribution was predicted with MRS(G) LQ PDF's). Both ratios are plotted in right panel of Fig. 2. Our result is systematically shifted down with respect to the experimental data.

---

[4] Gauge invariance implies that as $F_2^{ep,res}(Q^2_{res} \to 0) \to 0$; as a consequence, in the ratio of Eq. 4, the numerator decreases faster than the denominator.

In Ref. [4] it was shown that if one assumes the BG duality for F_2^{ep} structure function, then the resonance data can be approximated by the DIS-like $F_2^{res-par}$ parametrization. The obtained fit (on an average level) describes well resonance data. We use this parametrization in our analysis. In Fig. 2 we plot the ratio

$$R_2^{res-par}\left(Q^2_{res}\right) = \frac{\displaystyle\int_{\xi_{min}}^{\xi_{max}} d\xi\, F_2^{ep,RS}(\xi, Q^2_{res})}{\displaystyle\int_{\xi_{min}}^{\xi_{max}} d\xi\, F_2^{res-par}(\xi)}, \qquad (5)$$

computed for $1.1 < W^2 < 3.9$ GeV^2. We see that the above function weakly depends on Q^2_{res}. Additionally, in almost all considered domains $R_2^{res-par}\left(Q^2_{res}\right)$ is around 0.6. It illustrates the fact that the quark model systematically underestimates the experimental data, and it needs to be enriched by the nonresonant background contribution. On the other hand this regular deviation between quark model predictions and the data allows us to propose the effective structure function as:

$$\begin{aligned} F^{ep}_{2,dual}(\xi) &= F_2^{ep,res}(\xi) + 0.4\eta(\xi) F_2^{ep,val}, \qquad (6) \\ \eta(\xi) &= 2\arctan((W - M - m_\pi)/GeV)/\pi, (7) \end{aligned}$$

where the $F_2^{ep,val}$ is the valence DIS contribution, while $\eta(\xi)$ function is introduced to make a smooth transition from the threshold to the $P_{33}(1232)$ resonance peak. It is clear the ratio $R\left(F^{ep}_{2,dual}, Q^2_{res}, Q^2_{DIS} = 10\ \text{GeV}^2\right)$ and the analogously computed ratio (Eq. 5) will tend to one. In Fig. 2 (left panel), we plot $F^{ep}_{2,dual}$ together with the CLAS data. The good agreement in the first and also second resonance region can be seen.

The above approximation of the structure function seems to be naive in simplicity but very effective in the applications. In particular, it can be useful to apply to the Monte Carlo generators, dedicated to simulate neutrino-matter interactions.

NEUTRINO-NUCLEON INTERACTIONS

It is well known that if the $P_{33}(1232)$ resonance region is included in the discussion then the BG duality can not be simultaneously observed for the proton and neutron targets. It is the result of the isospin symmetry of the resonance structure function and $SU(6)$ symmetry property of the DIS structure function. Fortunately the BG duality (assuming some accuracy) appears for the weak structure functions of the isoscalar target .

For the study of νN scattering we apply the updated RS model but the description is supplemented by including the nonresonant background contribution (we follow the effective description proposed in Ref. [12]).

We consider the weak charged current (CC) and neutral current (NC) nucleon structure functions. The following ratios are computed:

$$\begin{aligned} R_1^{\nu N}(Q^2_{res}) &\equiv R\left(xF_1^{\nu N}, Q^2_{res}, Q^2_{DIS} = 10\ \text{GeV}^2\right), \quad (8) \\ R_2^{\nu N}(Q^2_{res}) &\equiv R\left(F_2^{\nu N}, Q^2_{res}, Q^2_{DIS} = 10\ \text{GeV}^2\right), \quad (9) \\ R_3^{\nu N}(Q^2_{res}) &\equiv R\left(xF_3^{\nu N}, Q^2_{res}, Q^2_{DIS} = 10\ \text{GeV}^2\right) \quad (10) \end{aligned}$$

We found out that for $W_{max} = 1.8$ GeV (see Fig. 3) the above ratios, similarly as for ep scattering, weakly depend on Q^2_{res}. Once again it can be treated as evidence for appearance of the duality.

It is interesting that the ratios of CC and NC structure functions are very similar. It is the result of the $SU(3) \times SU(2)$ and $SU(6)$ symmetries of the structure functions. As a consequence if the duality is observed for the one channel, say CC, it must be also visible (on the same level) for the NC structure functions. This property can have applications in modeling of the cross sections for the neutral current neutrino-nucleus scattering.

As was mentioned in first section, the predictions of the cross sections for neutrino-nucleon scattering are affected by the lack of knowledge of the axial structure of the nucleon. Therefore we present the ratios (8-10) with 1σ error coming from the axial form factor uncertainty (we apply the results of Ref. [19], where the uncertainty of the axial contribution was discussed). It is seen (in Fig. 3) that for the $xF_3^{\nu N}$ and $xF_1^{\nu N}$ structure functions the BG duality is observed within 1σ error, whereas for $F_2^{\nu N}$ structure function the duality appears within 2σ error.

However, we remark that in the discussion of structure function uncertainties, one needs to remember that in our approach the description of NoBG contribution is quite approximate. It gives rise to a systematic error. In order to show the role of NoBG contribution we plot the ratios (Eqs. 8–10) computed without NoBG.

OUTLOOK

As was mentioned in the first section, the duality is observed in electron-deuteron scattering. But the scaling behavior of the resonance data for electron scattering off heavier nuclear targets, like carbon and iron [20], is also observed. Here the duality between the resonance and the DIS nuclear structure function is visible even before averaging the data.

In fact, the Fermi motion effect naturally ”averages” the nucleon structure function over a wide kinematical region, but, there are also other nuclear medium effects like renormalization of the resonance properties, redistribution of the resonance prominent regions etc.. It makes the theoretical investigation of the duality in lepton scattering off nuclei very difficult and delicate.



FIGURE 3. Plots of Eqs. 8, 9, and 10, computed with $W_{max} = 1.8$ GeV are shown. The solid lines denote the full model predictions with NoBG. The computations without NoBG contribution are denoted by dashed lines. The thick and thin lines denote the ratios for CC and NC structure functions. The shadow areas denote the 1σ uncertainty.

On the other hand, observing the duality for nuclear targets means that the nuclear corrections in the resonance and the DIS regions are comparable. But it has been not expected [2].

Some effort to investigate the duality in lepton scattering off nuclei was done by Lalakulich *et al.* [21]. The electromagnetic and weak nuclear structure functions were obtained with an independent shell particle model, and then compared with the experimental data. For the ν-nuclei scattering the duality was not obtained. However, probably further studies of the nuclear effects and the role of NoBG contribution can shed light on this topic [22].

It is obvious that assuming that the Bloom-Gilman duality is also a fundamental property of weak nuclear structure functions can give interesting applications in neutrino scattering physics. In particular in can be useful for:

(i) modeling the neutrino cross sections in the resonance region;
(ii) extraction of the axial contribution from the scattering data;
(iii) studying the nonresonant background contribution.

Hopefully the forthcoming neutrino experiments, like Minerνa will be able to critically investigate the duality in neutrino scattering off nuclei.

ACKNOWLEDGMENTS

The author was supported by the grant: 35/N-T2K/2007/0 (the project number DWM/57/T2K/2007).

REFERENCES

1. E. D. Bloom and F. J. Gilman, *Phys. Rev. Lett.* **25**, 1140 (1970); *Phys. Rev.* **D4**, 2901 (1971).
2. W. Melnitchouk *et al.*, *Phys. Rept.* **406**, 127 (2005).
3. A. De Rujula *et al.*, *Annals Phys.* **103**, 315 (1977).
4. I. Niculescu *et al.*, *Phys. Rev. Lett.* **85**, 1186 (2000).
5. I. Niculescu *et al.*, *Phys. Rev. Lett.* **85**, 1182 (2000).
6. J. Morfin, *PDFs from neutrino DIS review*, the talk given at NuFact09, 11th International Workshop on Neutrino Factories, Superbeams and Beta Beams, July 20-25, 2009 – Illinois Institute of Technology – Chicago, USA.
7. K. Matsui *et al.*, *Phys. Rev.* **C72**, 025204 (2005).
8. K. M. Graczyk *et al.*, *Nucl. Phys.* **A781**, 227 (2007).
9. O. Lalakulich *et al.*, *Phys. Rev.* **C75**, 015202 (2007).
10. A. Bodek *et al.*, *Eur. Phys. J.* **C53** (2008) 349.
11. K. M. Graczyk *et al.*, *Phys. Rev.* **D77**, 053001 (2008) [Erratum-ibid. **D79**, 079903(E) (2009)], ibid. **D77**, 053003 (2008).
12. D. Rein, L.M. Sehgal, *Ann. Phys.* **133**, 79 (1981).
13. R.P. Feynman, *et al.*, *Phys. Rev.* **D3**, 2706 (1971); F. Ravndal, Lett. Nuovo Cimento, **3**, 631 (1972) and *Phys. Rev.* **D4**, 1466 (1971).
14. J. A. Nowak [MiniBooNE Collaboration], arXiv:0909.3659.
15. F. E. Close, *at al.*, *Phys. Rev.* **D6**, 2533 (1972); F. E. Close and F. J. Gilman, *Phys. Rev.* **D7**, 2258 (1973); F. E. Close and W. Melnitchouk, *Phys. Rev.* **C68**, 035210 (2003).
16. M. Glück, *et al.*, *Z. Phys.* **C67**, 433 (1995).
17. F. E. Close and N. Isgur, *Phys. Lett.* **B509**, 81 (2001).
18. M. Osipenko *et al.* [CLAS Collaboration], *Phys. Rev.* **D67** (2003) 092001. M. Osipenko et al., *The proton sructure function F(2) with CLAS.* CLAS-NOTE-2003-001 (Sep 2003) arXiv:0309.052.
19. K. M. Graczyk, D. Kielczewska, P. Przewlocki and J. T. Sobczyk, Phys. Rev. D **80**, 093001 (2009)
20. J. Arrington *et al.*, *Phys. Rev.* **C73**, 035205 (2006).
21. O. Lalakulich *et al.*, *Phys. Rev.* **C79**, 015206 (2009).
22. T. Leitner, O. Lalakulich, O. Buss, U. Mosel and L. Alvarez-Ruso, arXiv:0910.2835 [nucl-th].